

## **Informed Consent – Poly lactic acid**

**Dr. Alessio Redaelli**

**Via Di Vittorio 24 20070 DRESANO**

### **Informed consent for implant of poly lactic acid (New Filler)**

I undersigned \_\_\_\_\_ age \_\_\_\_\_  
living in \_\_\_\_\_ street \_\_\_\_\_  
phone number \_\_\_\_\_ e-mail \_\_\_\_\_

In view of the medical-therapeutic act above indicated, I confirm that I have been well informed about the nature and the known effects of the medical operation described later. The consent is personal and it cannot be delegated to the family (if not for minor or under guardianship).

Poly lactic acid is a polymer of the lactic acid, it is produced by chemical synthesis and it has a totally biodegradability and biocompatibility if injected in human tissues.

This substance is exempt from risk of allergic reactions (as it results from a wide literature on a world level). With poly lactic acid is possible to reduce the depth of wrinkles and increase the subcutaneous volumes.

The immediate effect is the fullness due to the quantity of injected liquid which the active principle was diluted with: it is fundamental to understand and be well informed on the temporariness of this result in order to avoid dissatisfaction. The real result, in fact, appears later on during the first 3-4 sessions (which means about 2-3 months later) and it will reach the maximum clinical evidence also after many months from the first implant. This time is necessary in order for the poly lactic acid to carry out the action of stimulating the production of new collagen.

In the 2-3 following days, it is important to avoid the exposure to very hot or very cold temperature.

The duration of the correction is variable in base of the frequency with which the zone of the implant is stimulated; basically the result reaches its maximum from 6-8 till 12-14 months after the first implant, but it varies from patient to patient.

The implant of the poly lactic acid is not particularly painful. An anesthetic lotion is used half an hour before the treatment. Sometime an injective anesthesia is required.

In literature very rare inflammatory and infective reactions occur, and rarely the appearance of long term subcutaneous nodules.

Sessions for additional touches may occur also even after short time from the operation.

The procedures and the used equipments are considered the most suitable from a scientific-medical point of view.

I declare however to be allergic to these following substances:

.....

and to have already done the following aesthetic therapies:

.....

After having taken note of the illustrated situation, I accept the suggested medical procedure.

Date

\_\_\_\_\_

Patient's Signature

\_\_\_\_\_

Doctor's Signature

\_\_\_\_\_